



























- Rasheed Z, Akhtar N, Haqqi TM. Advanced glycation end products induce the expression of interleukin-6 and interleukin-8 by receptor for advanced glycation end product-mediated activation of mitogen-activated protein kinases and nuclear factor- $\kappa$ B in human osteoarthritis chondrocytes. *Rheumatology (Oxford)* 2011; 50(5): 838-51.
- Singh VP, Bali A, Singh N, Jaggi AS. Advanced glycation end products and diabetic complications. *Korean J Physiol Pharmacol* 2014; 18(1): 1-14.
- Stevens AL, Wishnok JS, White FM, Grodzinsky AJ, Tannenbaum SR. Mechanical injury and cytokines cause loss of cartilage integrity and upregulate proteins associated with catabolism, immunity, inflammation, and repair. *Mol Cell Proteomics* 2009; 8(7): 1475-89.
- Syggelos SA, Aletras AJ, Smirlaki I, Skandalis SS. Extracellular matrix degradation and tissue remodeling in periprosthetic loosening and osteolysis: focus on matrix metalloproteinases, their endogenous tissue inhibitors, and the proteasome. *Biomed Res Int* 2013; 2013: 230805.
- Thai Health Promotion Foundation. NCDs (Non-Communicable diseases) [online] 2017 [cited 2017 Dec 11]. Available from: <http://www.thaihealth.or.th>
- Verzijl N, DeGroot J, Thorpe SR, Bank RA, Shaw JN, Lyons TJ, et al. Effect of collagen turnover on the accumulation of advanced glycation end products. *J Biol Chem* 2000; 275(50): 39027-31.
- World Health Organization. Complications of diabetes [online] 2014 [cited 2018 Jun 4]. Available from: [https://www.who.int/diabetes/action\\_online/basics/en/index3.html](https://www.who.int/diabetes/action_online/basics/en/index3.html)
- World Health Organization. Noncommunicable Diseases Progress Monitor 2017 [online] 2017 [cited 2017 Dec 24]. Available from: <https://www.who.int/nmh/publications/ncd-progress-monitor-2017/en/>